

Community pharmacist management of discharge medication summaries in primary care

Michael Wilcock,¹ *Associate Editor*, David Bearman,² *Clinical lead*

¹*Department of Pharmacy, Royal Cornwall Hospitals NHS Trust, Truro, UK*

²*Peninsula Academic Health Science Network, Exeter, UK*

Correspondence to Mr Michael Wilcock; mike.wilcock@nhs.net

Transition of care, when a patient moves from one setting of care to another, is a period of high vulnerability for patients and one of the six most common areas of risk with medicines across health and adult social care services.^{1–4} For example, after discharge from hospital, problems may arise when patients are not prescribed their usual medicines. Other known issues include transcribing errors, mistakes or omissions in discharge letters issued by hospitals to general practitioners (GPs), as well as delays in acting on the information provided in the discharge summary.^{3–5} This can result in patients moving between care settings without a full understanding of any new medicines or changes to their existing medicines.

In general, community pharmacists are not routinely sent information on discharge medication for patients who have been in hospital, or even informed that patients have been admitted or discharged. However, evidence is emerging that involving community pharmacists in the process of medicines optimisation after hospital discharge can improve medicines reconciliation and patient safety, and may be associated with shorter hospital stays and lower readmission rates.^{6–11} These reports of patient benefit resulting from communication with, and involvement of, the community pharmacist at discharge have appeared from different countries with different health systems, using different community pharmacist interventions.^{11–14} A pilot scheme in the UK has shown that it is possible to transfer discharge information electronically from secondary care to allow community pharmacies to provide a follow-up consultation tailored to the patient's need.¹⁰ It also indicated that patients may have lower rates of readmission and shorter hospital stays. Evaluation of the Discharge Medicines

Review Service provided by community pharmacists in Wales suggested that the scheme was cost saving through reduced accident and emergency attendances, hospital admissions and drug wastage. Of more importance were the health benefits to patients that resulted from the avoidance of adverse events.¹⁵

Nevertheless, these and other unpublished UK-based studies of transfer of care around medicines (TCAM) projects have varied both in the electronic referral platforms used and how the service is delivered. For example, different patient cohorts were used as target groups, and there were different approaches to the use of telephone Medicines Use Reviews as an intervention offered by the community pharmacist. These UK evaluations are also not consistent in the framework they used or the data they collected, which makes firm analysis of the benefits problematic. Despite these uncertainties, there is the prospect that such schemes could help reduce the pressure on secondary care services and release value to the health system.

However, current performance across hospital trusts participating in TCAM services shows variable levels of both referral and community pharmacy completion rates (offering considerable scope for improvement). As yet, there is no structured information of the benefit levels that are likely to be achieved in different patient cohorts, and therefore the scale of the potential benefit offered by TCAM, while large, remains unknown. To improve the current performance, key areas need to be addressed. These include an understanding of which patient cohorts benefit the most; a support process in secondary care to enhance the referral rate (eg, improved technology that fits easily into the hospital pharmacy team workflow); and targeted investment within the community pharmacy contractual framework to deliver medicines

optimisation after hospital discharge. There should also be an oversight function to monitor and support completion of referrals as well as evaluation and feedback to support quality improvement.

At the same time, it is important to recognise that another approach to the management of discharge summaries has emerged through the introduction of clinical pharmacists in general practice in England.¹⁶ Evaluation of the pilot scheme identified that a large proportion of pharmacists were responsible for streamlining the discharge reconciliation process for patients. It was suggested that such work had a significant positive impact on patients and had the potential to reduce hospital readmissions. These findings need to be backed up by robust evidence of benefit to patients. A coordinated approach to the management of discharge summaries by community pharmacists and pharmacists working in general practice will be needed to avoid duplication of effort.

Although TCAM is one of the Academic Health Science Network priorities for national adoption and spread, it will need support from commissioners, GP practices, pharmacies and patients that a clinical handover service from hospitals to pharmacists in the community is wanted by all stakeholders.¹⁷ Such a service should clearly define who does what for whom and when.

Competing interests None declared. Disclosure of conflicts of interest form(s) are published online as supplementary files.

Provenance and peer review Commissioned; externally peer reviewed.

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

References

- 1 World Health Organization. *Medication without harm - global patient safety challenge on medication safety* [online]. 2017. Available: <https://www.who.int/patientsafety/medication-safety/en/> [Accessed 29 August 2019].
- 2 World Health Organization. *Medication safety in transitions of care* [online]. 2019. Available: <https://apps.who.int/iris/bitstream/handle/10665/325453/WHO-UHC-SDS-2019.9-eng.pdf> [Accessed 29 August 2019].
- 3 Care Quality Commission. *Medicines in health and adult social care* [online]. 2019. Available: https://www.cqc.org.uk/sites/default/files/20190605_medicines_in_health_and_adult_social_care_report.pdf [Accessed 29 August 2019].
- 4 Kripalani S, LeFevre F, Phillips CO, et al. Deficits in communication and information transfer between hospital-based and primary care physicians - implications for patient safety and continuity of care. *J Am Med Assoc* 2007;297:831–41.
- 5 Shah C, Hough J, Jani Y. Medicines reconciliation in primary care: a study evaluating the quality of medication-related information provided on discharge from secondary care. *Eur J Hosp Pharm* 2018;0:1–6.
- 6 Paulino EI, Bouvy ML, Gastelurrutia MA, et al. Drug related problems identified by European community pharmacists in patients discharged from hospital. *Pharm World Sci* 2004;26:353–60.
- 7 Duggan C, Feldman R, Hough J, et al. Reducing adverse prescribing discrepancies following hospital discharge. *Int J Pharm Pract* 1998;6:77–82.
- 8 Hockly M, Williams S, Allen M, et al. Transfer of care - a randomised control trial investigating the effect of sending the details of patients' discharge medication to their community pharmacist on discharge from hospital. *Int J Pharm Pract* 2018;26:174–82.
- 9 Nazar H, Nazar Z, Portlock J, et al. A systematic review of the role of community pharmacies in improving the transition from secondary to primary care. *Br J Clin Pharmacol* 2015;80:936–48.
- 10 Nazar H, Brice S, Akhter N, et al. New transfer of care initiative of electronic referral from hospital to community pharmacy in England: a formative service evaluation. *BMJ Open* 2016;6:e012532.
- 11 Sabir FRN, Tomlinson J, Strickland-Hodge B, et al. Evaluating the connect with pharmacy web-based intervention to reduce Hospital readmission for older people. *Int J Clin Pharm* 2019.
- 12 Daliri S, Hugtenburg JG, Ter Riet G, et al. The effect of a pharmacy-led transitional care program on medication-related problems post-discharge: a before-after prospective study. *PLoS One* 2019;14:e0213593.
- 13 Ensing HT, Koster ES, Dubero DJ, et al. Collaboration between hospital and community pharmacists to address drug-related problems: the HomeCoMe-program. *Res Social Adm Pharm* 2019;15:267–78.
- 14 Shaver A, Morano M, Pogodzinski J, et al. Impact of a community pharmacy transitions-of-care program on 30-day readmission. *J Am Pharm Assoc* 2019;59:202–9.
- 15 Welsh Institute for Health and Social Care. *Evaluation of the discharge medicines review service* [online]. 2014. Available: [http://www.cpwales.org.uk/Contract-support-and-IT/Advanced-Services/Discharge-Medicines-Review-\(DMR\)/Evaluation-of-the-DMR-Service/Evaluation-of-the-DMR-service.aspx](http://www.cpwales.org.uk/Contract-support-and-IT/Advanced-Services/Discharge-Medicines-Review-(DMR)/Evaluation-of-the-DMR-Service/Evaluation-of-the-DMR-service.aspx) [Accessed 29 August 2019].
- 16 Mann C, Anderson C, Avery AJ, et al. *Clinical pharmacists in General Practice: pilot scheme independent evaluation report: full report* [online]. 2018. Available: <https://www.nottingham.ac.uk/pharmacy/documents/generalpracticeyearfwdrev/clinical-pharmacists-in-general-practice-pilot-scheme-full-report.pdf> [Accessed 29 August 2019].
- 17 The AHSN network. *Transfers of care around medicines (TCAM)* [online]. 2018. Available: <https://www.ahsnnetwork.com/about-academic-health-science-networks/national-programmes-priorities/transfers-care-around-medicines-tcam> [Accessed 29 August 2019].

DOI: 10.1136/dtb.2019.000052